

Figure 1





Figure 2

Betting Log - Type 2 sports

Figure 3



Betting Log Type 1 sports

Figure 4





FIGURE 5. Penalties for choosing  $f=f_e \neq f^*=f_t$



## APPENDIX I. Integrals for deriving moments of $E_\infty$

$$I_0(a^2, b^2) = \int_0^\infty \exp[-(a^2x^2 + b^2/x^2)] dx$$

$$I_n(a^2, b^2) = \int_0^\infty x^n \exp[-(a^2x^2 + b^2/x^2)] dx$$

Given  $I_0$  find  $I_2$

$$I_0(a^2, b^2) = \int_0^\infty \exp[-(a^2x^2 + b^2/x^2)] dx$$

$$= - \int_\infty^0 \exp[-(a^2/u^2 + b^2u^2)] (-du/u^2)$$

where  $x = 1/u$  and  $dx = -du/u^2$  so

$$I_0(a^2, b^2) = \int_0^\infty x^{-2} \exp[-(b^2x^2 + a^2/x^2)]$$

$$= I_{-2}(b^2, a^2)$$

$$(A1) \quad \text{hence } I_{-2}(a^2, b^2) = I_0(b^2, a^2) = \frac{\sqrt{\pi}}{2|b|} e^{-2|ab|}$$

$$I_0 = \int_0^\infty \exp[-(a^2x^2 + b^2/x^2)] dx = U \cdot V \Big|_0^\infty - \int_0^\infty V dU$$

where  $U = \exp[\cdot]$ ,  $dV = dx$ ,  $dU = (\exp[\cdot])(-2a^2x + 2b^2x^{-3})$  and  $V = x$  so

$$I_0 = \exp[-(a^2x^2 + b^2/x^2)] \cdot x \Big|_0^\infty - \int_0^\infty (-2a^2x^2 + 2b^2/x^2) \exp[-(a^2x^2 + b^2/x^2)] dx$$

$$= 2a^2 I_2(a^2, b^2) - 2b^2 I_{-2}(a^2, b^2)$$

Hence:

$$I_0(a^2, b^2) = 2a^2 I_2(a^2, b^2) - 2b^2 I_{-2}(a^2, b^2)$$

and  $I_{-2}(a^2, b^2) = I_0(b^2, a^2)$  so substituting and solving for  $I_2$  gives

$$I_2(a^2, b^2) = \frac{1}{2a^2} \{I_0(a^2, b^2) + 2b^2 I_0(b^2, a^2)\}$$

Comments.

1. We can solve for all even  $n$  by using  $I_0, I_{-2}$  and  $I_2$ , and integration by parts.
2. We can use the indefinite integral  $J_0$  corresponding to  $I_0$ , and the previous methods, to solve for  $J_{-2}, J_2$ , and then for all even  $n$ . Since

$$\begin{aligned} I_0(a^2, b^2) &= \frac{\sqrt{\pi}}{2|a|} e^{-2|ab|} && \text{then} \\ I_{-2}(b^2, a^2) &= \frac{\sqrt{\pi}}{2|b|} e^{-2|ab|} && \text{and} \\ I_2(a^2, b^2) &= \frac{1}{2a^2} \left\{ \frac{\sqrt{\pi}}{2|a|} + 2b^2 \frac{\sqrt{\pi}}{2|b|} \right\} e^{-2|ab|} \\ &= \frac{\sqrt{\pi}}{4a^2} e^{-2|ab|} \{1/|a| + 2|b|\} \end{aligned}$$

## APPENDIX II. Derivation of Formula (3.1)

This is based on a note from Howard Tucker. Any errors are mine.

\*From the paper by Paranjape & Park, if  $X(t)$  is standard Brownian motion, if  $a \neq 0$ ,  $b > 0$ ,

$$\begin{aligned} & P(X(t) \leq at + b, 0 \leq t \leq T \mid X(T) = s) \\ &= \begin{cases} 1 - \exp\left\{-\frac{2b}{T}(aT + B - s)\right\} & \text{if } s \leq aT + b \\ 0 & \text{if } s > aT + B. \end{cases} \end{aligned}$$

Write this as:

$$\begin{aligned} & P(X(t) \leq at + b, 0 \leq t \leq T \mid X(T)) \stackrel{a.s.}{=} \\ &= (1 - \exp\{-2b(aT + b - X(T))\frac{1}{T}\}) \quad \text{if } X(T) \leq aT + b. \end{aligned}$$

Taking expectations of both sides of the above, we get

$$\begin{aligned} & P(X(t) \leq at + b, 0 \leq t \leq T) \\ &= \int_{-\infty}^{aT+b} \left(1 - e^{-2b(aT+b-s)/T}\right) \frac{1}{\sqrt{2\pi T}} e^{-s^2/2T} ds \\ &= \frac{1}{\sqrt{2\pi T}} \int_{-\infty}^{aT+b} e^{-s^2/2T} ds - \frac{e^{-2ab}}{\sqrt{2\pi T}} \int_{-\infty}^{aT+b} e^{-(s-2b)^2/2T} ds \end{aligned}$$

Hence

(I)  $P(X \text{ going above line } at + b \text{ during } [0, T]) = 1 - \text{previous probability}$

$$= \frac{1}{\sqrt{2\pi T}} \int_{aT+b}^{\infty} e^{-s^2/2T} ds + e^{-2ab} \cdot \frac{1}{\sqrt{2\pi T}} \int_{-\infty}^{aT-b} e^{-u^2/2T} du, \text{ where } u = s - 2b.$$

Now, when  $a = 0$ ,  $b > 0$ ,

$$P\left[\sup_{0 \leq t \leq T} X(t) \geq b\right] = \sqrt{\frac{2}{\pi T}} \int_b^{\infty} e^{-v^2/2T} dv,$$



which agrees with a known formula (see, e.g., page 261 of Tucker [1967]). When  $T \rightarrow \infty$ , since  $\sqrt{T}/T \rightarrow 0$  and  $\sqrt{T} = s.d. \text{ of } X(T)$ , the first integral  $\rightarrow 0$ , the second integral  $\rightarrow 1$ , and  $P(X \text{ ever rises above line } at + b) = e^{-2ab}$ .

\*The theorem it comes from is due to Sten Malmquist: On Certain Confidence Countours for Distribution Functions, Ann. Math. Stat., 25 (1954), pp 523-533. This theorem is stated in S.R. Paranjape and C. Park: Distribution of the Supremum of the Two-Parameter Yeh-Wiener Process on the Boundary, J. Appl. Prob. Vol. 10 (1973).

Letting  $\alpha = a\sqrt{T}$ ,  $\beta = b/\sqrt{T}$ , formula I becomes

$$P(\cdot) = N(-\alpha - \beta) + e^{-2\alpha\beta}N(\alpha - \beta) \text{ where } \alpha, \beta > 0 \text{ or}$$

$$(II) \quad P(X(t) \leq at + b, 0 \leq t \leq T) = 1 - P(\cdot) = N(\alpha + \beta) - e^{-2\alpha\beta}N(\alpha - \beta)$$

for the probability the line is never surpassed. This follows from:

$$\frac{1}{\sqrt{2\pi T}} \int_{aT+b}^{\infty} e^{-s^2/2T} ds = \frac{1}{\sqrt{2\pi}} \int_{a\sqrt{T}+b/\sqrt{T}}^{\infty} e^{-x^2/2} dx = N(-\alpha - \beta)$$

$$\text{and} \quad \frac{1}{\sqrt{2\pi T}} \int_{-\infty}^{aT-b} e^{-u^2/2T} du = N(\alpha - \beta)$$

where  $s = aT + b$  and  $x = s/\sqrt{T} = a\sqrt{T} + b/\sqrt{T}$   
and  $\alpha = a\sqrt{T}$  and  $\beta = b/\sqrt{T}$ .

The formula becomes:

$$P(\sup[X(t) - (at + b)] \geq 0 : 0 \leq t \leq T)$$

$$\begin{aligned} &= N(-\alpha - \beta) + e^{-2\alpha\beta}N(\alpha - \beta) \\ &= N(-\alpha - \beta) + e^{-2\alpha\beta}N(\alpha - \beta) \quad \alpha, \beta > 0 \end{aligned}$$

Observe that  $P(\cdot) < N(-\alpha - \beta) + N(\alpha - \beta)$

$$\begin{aligned} &= \{1 - N(\alpha + \beta)\} + N(\alpha - \beta) \\ &= \int_{-\infty}^{\alpha-\beta} \alpha(x) dx + \int_{\alpha+\beta}^{\infty} \alpha(x) dx < 1 \end{aligned}$$

as it should be.

## Appendix III. Expected time to reach goal

Reference: Handbook of Mathematical Functions, Abramowitz and Stegun, Editors, N.B.S. Applied Math. Series 55, June 1964.

p.304, 7.4.33 gives with  $erfz \equiv \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt$  the integral:

$$(1) \quad \int \exp\{-(a^2x^2 + b^2/x^2)\} dx = \frac{\sqrt{\pi}}{4a} \left[ e^{2ab} erf(ax + b/x) + e^{-2ab} erf(ax - b/x) \right] + C,$$

$a \neq 0$ .

Now the left side is  $> 0$  so for real  $a$ , we require  $a > 0$  otherwise the right side is  $< 0$ , a contradiction.

We also note that p.302, 7.4.3. gives

$$(2) \quad \int_0^\infty \exp\{-(at^2 + b/t^2)\} dt = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{-2\sqrt{ab}}$$

with  $\Re a > 0$ ,  $\Re b > 0$ .

To check (2) v. (1), suppose in (1)  $a > 0$ ,  $b > 0$  and find

$\lim_{x \rightarrow 0} \text{ and } \lim_{x \rightarrow \infty}$  of  $erf(ax + b/x)$  and  $erf(ax - b/x)$

$$\lim_{x \downarrow 0+} (ax + b/x) = +\infty \quad \lim_{x \downarrow 0+} (ax - b/x) = -\infty$$

$$\lim_{x \rightarrow \infty} (ax + b/x) = +\infty \quad \lim_{x \rightarrow \infty} (ax - b/x) = +\infty$$

(1) becomes

$$\begin{aligned} \frac{\sqrt{\pi}}{4a} e^{-2ab} [erf(\infty) - erf(-\infty)] &= \frac{\sqrt{\pi}}{4a} e^{-2ab} 2erf(\infty) \\ &= \frac{\sqrt{\pi}}{2a} e^{-2ab} \end{aligned}$$

since we know  $erf(\infty) = 1$ .

In (2) replace  $a$  by  $a^2$ ,  $b$  by  $b^2$  to get

$$I_0(a^2, b^2) \equiv \int_0^\infty \exp\{-(a^2t^2 + b^2/t^2)\} dt = \frac{1}{2} \frac{\sqrt{\pi}}{|a|} e^{-2|ab|}$$

which is the same.

Note: if we choose the lower limit of integration to be 0 in (1), then we can find  $C$ :

$$\begin{aligned}
 0 &= \int_0^0 \exp\{-(a^2x^2 + b^2/x^2)\}dx \\
 &= \frac{\sqrt{\pi}}{4a} [e^{2ab} \operatorname{erf}(\infty) + e^{-2ab} \operatorname{erf}(-\infty)] + C \\
 &= \frac{\sqrt{\pi}}{4a} [e^{2ab} - e^{-2ab}] + C.
 \end{aligned}$$

Whence

$$\begin{aligned}
 (3) \quad F(x) &\equiv \int_0^x \exp\{-(a^2x^2 + b^2/x^2)\}dx \\
 &= \frac{\sqrt{\pi}}{4a} \{e^{2ab} [\operatorname{erf}(ax + b/x) - 1] + e^{-2ab} [\operatorname{erf}(ax - b/x) + 1]\}
 \end{aligned}$$

To see how (3) might have been discovered, differentiate:

$$\begin{aligned}
 F'(x) &= \exp\{-(a^2x^2 + b^2/x^2)\} \\
 &= \frac{\sqrt{\pi}}{4a} \{e^{2ab}(a - b/x^2) \operatorname{erf}'(ax + b/x) + e^{-2ab}(a + b/x^2) \operatorname{erf}'(ax - b/x)\}
 \end{aligned}$$

Now  $\operatorname{erf}'(z) = \frac{2}{\sqrt{\pi}} \exp(-z^2)$  so

$$\begin{aligned}
 \operatorname{erf}'(ax + b/x) &= \frac{2}{\sqrt{\pi}} \exp[-(ax + b/x)^2] \\
 &= \frac{2}{\sqrt{\pi}} \exp\{-(a^2x^2 + b^2/x^2 + 2ab)\} \\
 &= \frac{2}{\sqrt{\pi}} e^{-2ab} \exp\{-(a^2x^2 + b^2/x^2)\}
 \end{aligned}$$

and, setting  $b \leftarrow -b$ ,

$$\operatorname{erf}'(ax - b/x) = \frac{2}{\sqrt{\pi}} e^{2ab} \exp\{-(a^2x^2 + b^2/x^2)\}$$

whence

$$\begin{aligned}
 F'(x) &= \frac{\sqrt{\pi}}{4a} \left\{ \frac{2}{\sqrt{\pi}} (a - b/x^2) + \frac{2}{\sqrt{\pi}} (a + b/x^2) \right\} \exp\{-(a^2x^2 + b^2/x^2)\} \\
 &= \frac{1}{2a} \{2a\} \exp\{-(a^2x^2 + b^2/x^2)\} \\
 &= \exp\{-(a^2x^2 + b^2/x^2)\}
 \end{aligned}$$

Case of interest:  $a < 0, b > 0$

Expect:

$b > 0, a \leq 0 \Rightarrow F(T) \uparrow 1 \text{ as } T \rightarrow \infty$

$b > 0, a > 0 \Rightarrow F(T) \uparrow c < 1 \text{ as } T \rightarrow \infty$

If  $b > 0, a = 0$ :

$$F(T) = N(-\beta) + N(-\beta) = 2N(-b/\sqrt{T}) \uparrow 2N(0) = 1 \text{ as } T \uparrow \infty.$$

Also, as expected  $F(T) \uparrow 1 \text{ as } b \downarrow 0$ .

If  $b > 0, a < 0$ : See Below.

If  $b > 0, a > 0$ :

$$\begin{aligned} F(T) &= N(-a\sqrt{T} - b/\sqrt{T}) + e^{-2ab}N(a\sqrt{T} - b/\sqrt{T}) \rightarrow N(-\infty) + e^{-2ab}N(\infty) \\ &= e^{-2ab} < 1 \text{ as } T \uparrow \infty. \quad \text{This is correct.} \end{aligned}$$

If  $b = 0$ :  $F(T) = N(-a\sqrt{T}) + N(a\sqrt{T}) = 1$ . This is correct.

Let  $F(T) = P(X(t) \geq at + b \text{ for some } t, 0 \leq t \leq T)$  which equals  $N(-\infty - \beta) + e^{-2ab}N(\infty - \beta)$  where  $\alpha = a\sqrt{T}$  and  $\beta = b/\sqrt{T}$  so  $ab = \alpha\beta$ ; we assume  $b > 0$  and  $a < 0$  in which case  $0 \leq F(T) \leq 1$  and  $\lim_{T \rightarrow \infty} F(T) = 1$ ,  $\lim_{T \rightarrow 0} F(T) = 0$ ;  $F$  is a probability distribution function:

$$\begin{aligned} \lim_{T \rightarrow 0} F(T) &= N(-\infty) + e^{-2ab}N(-\infty) = 0 \\ \lim_{T \rightarrow \infty} F(T) &= N(+\infty) + e^{-2ab}N(-\infty) = 1. \end{aligned}$$

The density function

$$\begin{aligned} f(T) &= F'(T) \\ &= \frac{\partial}{\partial T}(-\alpha - \beta)N'(-\alpha - \beta) + e^{-2ab} \frac{\partial}{\partial T}(\alpha - \beta)N'(\alpha - \beta) \end{aligned}$$

where

$$\frac{\partial \alpha}{\partial T} = \frac{1}{2}aT^{-1/2} \quad \frac{\partial \beta}{\partial T} = -\frac{1}{2}bT^{-3/2}$$

$$\begin{aligned} N'(-\alpha - \beta) &= \frac{1}{\sqrt{2\pi}} e^{-(\alpha+\beta)^2/2} = \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{(a^2T + b^2/T + 2ab)}{2} \right\} \\ N'(\alpha - \beta) &= \frac{1}{\sqrt{2\pi}} e^{-(\alpha-\beta)^2/2} = \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{(a^2T + b^2/T - 2ab)}{2} \right\} \end{aligned}$$

$$\begin{aligned}
Tf(T) &= T \left( -\frac{1}{2}aT^{-1/2} + \frac{1}{2}bT^{-3/2} \right) \frac{1}{\sqrt{2\pi}} e^{-ab} \exp \left\{ -\frac{(a^2T + b^2/T)}{2} \right\} \\
&+ Te^{-2ab} \left( \frac{1}{2}aT^{-1/2} + \frac{1}{2}bT^{-3/2} \right) \frac{1}{\sqrt{2\pi}} e^{ab} \exp \left\{ -\frac{(a^2T + b^2/T)}{2} \right\} \\
&= \frac{e^{-ab}}{2\sqrt{2\pi}} \left[ (-aT^{1/2} + bT^{-1/2}) \exp \left\{ -\frac{(a^2T + b^2/T)}{2} \right\} \right. \\
&\quad \left. + (aT^{1/2} + bT^{-1/2}) \exp \left\{ -\frac{(a^2T + b^2/T)}{2} \right\} \right]
\end{aligned}$$

The expected time to the goal is

$$\begin{aligned}
E_\infty = \int_0^\infty Tf(T)dT &= \frac{be^{-ab}}{\sqrt{2\pi}} \int_0^\infty T^{-1/2} \exp \left\{ -\frac{(a^2T + b^2/T)}{2} \right\} dT \\
\left. \begin{array}{l} T^{1/2} = x \\ T = x^2 \\ dT = 2xdx \end{array} \right\} &= \frac{2be^{-ab}}{\sqrt{2\pi}} \int_0^\infty \exp \left\{ -\left[ \left( \frac{a}{\pi^2} \right)^2 x^2 + \left( \frac{b}{\pi^2} \right)^2 x^{-2} \right] \right\} dx \\
&= \frac{2be^{-ab}}{\sqrt{2\pi}} I_0 \left( \left( \frac{a}{\sqrt{2}} \right)^2, \left( \frac{b}{\sqrt{2}} \right)^2 \right)
\end{aligned}$$

Now

$$\begin{aligned}
I_0(a^2, b^2) &= \frac{\sqrt{\pi}}{2|a|} e^{-2|ab|} \quad \text{so} \\
I_0 \left( \left( \frac{a}{\sqrt{2}} \right)^2, \left( \frac{b}{\sqrt{2}} \right)^2 \right) &= \frac{\sqrt{\pi}}{\sqrt{2}|a|} e^{-|ab|} \quad \text{whence} \\
E_\infty &= \frac{2be^{-ab}}{\sqrt{2\pi}} \frac{\sqrt{\pi}}{\sqrt{2}|a|} e^{-|ab|} = \frac{b}{|a|}, \quad a < 0, b > 0.
\end{aligned}$$

Note:

$$F'(T) = \frac{be^{-ab}}{\sqrt{2\pi}} T^{-3/2} \exp \left\{ -\frac{(a^2T + b^2/T)}{2} \right\} > 0$$

for all  $a$ , e.g.  $a < 0$ , so  $F(T)$  is monotone increasing. Hence, since  $\lim_{T \rightarrow \infty} F(T) = 1$ ,  $0 \leq F(T) \leq 1$  for all  $T$  so we have more confidence in using the formula for  $a < 0$  too.

$$\begin{array}{lll}
\text{Check:} & E_\infty(a, b) \downarrow 0 \text{ as } \downarrow -\infty & \text{yes} \\
& E_\infty(a, b) \uparrow \text{ as } b \uparrow & \text{yes}
\end{array}$$

$$\begin{array}{lll} E_{\infty}(a, b) \uparrow \text{ as } |a| \downarrow & \text{yes} \\ \text{note } \lim_{a \downarrow 0^+} E_{\infty}(a, b) = +\infty & \text{as suspected} \end{array}$$

This leads us to believe that in a fair coin toss (fair means no drift) and a gambler with finite capital, the expected time to ruin is infinite.

This is correct. Feller give  $D = z(a-z)$  as the duration of the game, where  $z$  is initial capital, ruin is at 0, and  $a$  is the goal. Then  $\lim_{a \rightarrow \infty} D(a) = +\infty$ .

NOTE:  $E_{\infty} = b/|a|$  means the expected time is the same as the point where  $aT + b$  crosses  $X(t) = 0$ . See Figure 2.

$$\begin{aligned} E_{\infty} &= b/|a| & a &= -m/s^2 & b &= \ln \lambda \\ \lambda &= C/X_0 = \text{normalized goal} \\ m &= p \ln(1+f) + q \ln(1-f) \equiv g(f) \\ s^2 &= pq \{ \ln[(1+f)/(1-f)] \}^2 \\ \text{Kelly fraction } f^* &= p - q & g(f^*) &= p \ln 2p + q \ln 2q \\ \text{For } m > 0, E_{\infty} &= (\ln \lambda) s^2 / g(f) \end{aligned}$$

Now this is the expected time in variance units. However  $s^2$  variance units = 1 trial so

$$n(\lambda, f) \equiv \frac{E_{\infty}}{s^2} = \frac{\ln \lambda}{g(f)} = \frac{\ln \lambda}{m}$$

is the expected number of trials.

$$\begin{aligned} \text{Checks: } n(\lambda, f) &\uparrow \text{ as } \lambda \uparrow \\ n(\lambda, f) &\rightarrow \infty \text{ as } \lambda \rightarrow \infty \\ n(\lambda, f) &\uparrow \text{ as } m \downarrow 0 \\ n(\lambda, f) &\rightarrow \infty \text{ as } m \rightarrow 0 \end{aligned}$$

Now  $g(f)$  has unique maximum at  $g(f^*)$  where  $f^* = p - q$ , the “Kelly fraction,” therefore  $n(\lambda, f)$  has a unique minimum for  $f = f^*$ . Hence  $f^*$  reaches a fixed goal in least expected time in this, the continuous case, so we must be asymptotically close to least expected time in the discrete case, which this approximates increasing by well in the sense of the CLT (Central Limit Theorem) and its special case, the normal approximation to the binomial distribution. The difference here is the trials are asymmetric. The positive and negative step sizes are unequal.

## References

- [1] Associates First Capital Corporation, *1997 Annual Report*. Associates First Capital Corporation, Dallas, Texas (1998).
- [2] Black, F. and M. Scholes, *The pricing of options and corporate liabilities.*, J. Polit. Econom., **81** (1973), 637-659.
- [3] Breiman, L., *Optimal gambling systems for favorable games.*, Fourth Berkeley Symposium on Probability and Statistics, **I**, (1961), 65-78.
- [4] Browne, S., *The Return on Investment From Proportional Portfolio Strategies*, (1995), Columbia University, New York.
- [5] Browne, S., *Reaching Goals by a Deadline: Continuous-Time Active Portfolio Management*, (1996), Columbia University, New York.
- [6] Buffett, Mary and Clark, David, *Buffettology*, Rawson Associates, Simon and Schuster, New York (1997).
- [7] Cox, D.R. and Miller, H.D., *The Theory of Stochastic Processes*, Wiley, New York (1965).
- [8] Feller, W., *An Introduction to Probability Theory and Its Applications*, Vol.I, Revised(1966). New York, John Wiley.
- [9] Griffin, P.A., *The Theory of Blackjack*, Las Vegas, Huntington Press, (1979); revised 1995.
- [10] Hagstrom, Robert G. Jr., *The Warren Buffett Way*, Wiley, New York (1994).
- [11] Hecht, R., private correspondence, (1995).
- [12] Ibbotson Associates, *1998 (or latest available) Yearbook: Stocks, Bonds, Bills and Inflation*, Ibbotson Associates, Chicago (1998).
- [13] Institutional Investor, Ivory Tower Investing, pages 43-55 (see page 44), March 1998.
- [14] Kelly, J.L., *A new interpretation of information rate*. Bell System Technical Journal, **35**, (1956), 917-926.

- [15] Kilpatrick, Andrew, *Of Permanent Value, the Story of Warren Buffet*. Distributed by Southern Publishers Group, Birmingham, Alabama (1994).
- [16] Kritzman, Mark, Risk and Utility: Basics, Chapter 2 of *Investment Management*, Bernstein and Damodaran, editors, Wiley, New York (1998).
- [17] Lowenstein, Roger, *The Making of an American Capitalist*. Random House, New York (1995).
- [18] Markowitz, H., *Portfolio Selection*, Cowles Monograph, **16**, Wiley, New York (1959).
- [19] McEnally, Richard W., *Latané's Bequest: The Best of Portfolio Strategies*. J. Portfolio Management, Vol. 12, No. 2, Winter 1986, 21-30.
- [20] Quaife, Art, *Rational Portfolio Determination*. The Trans Times, Vol. 2, No. 4, August 1993, Trans Time Inc., Oakland, California.
- [21] Quaife, Art, *Using the Sharpe Ratio to Evaluate Investments*. The Trans Times, Vol. 4, No. 1, February 1995, Trans Time Inc., Oakland, California.
- [22] Rotando, L. M. and E.O. Thorp, *The Kelly criterion and the stock market*. American Mathematical Monthly, (1992), 922-931.
- [23] Thorp, E.O., *Beat The Dealer*, Random House, New York (1962).
- [24] Thorp, E.O., *Beat The Dealer*, 2nd Ed., Vintage, New York (1966).
- [25] Thorp, E.O., *Optimal gambling systems for favorable games*. Review of the International Statistical Institute, **37**, (1969), 273-293.
- [26] Thorp, E.O., *Portfolio choice and the Kelly criterion*. Proceedings of the 1971 Business and Economics Section of the American Statistical Association, (1971), 215-224.
- [27] Thorp, E.O., *The Mathematics of Gambling*. Lyle Stuart, Secaucus, N.J., (1984).



- [28] Thorp, E.O. and Kassouf, S.T., *Beat The Market*, Random House, New York (1967).
- [29] Thorp, E.O. and Walden, W., *A winning bet in Nevada baccarat*. J. Amer. Statist. Assoc., **61**, Part I, (1966), 313-328.
- [30] Tucker, H., *A Graduate Course in Probability*, Academic Press, (1967).
- [31] Wong, S., *Professional Blackjack*, Pi Yee Press, (1994 edition), La Jolla, CA.